







Okeanos Explorer ROV Dive Summary

Dive Information	
Dive Map	The map displays the bathymetry of the Carondelet Reef area. Dive locations for EX-17-03 are marked with white circles. The map also shows the boundaries of the Howland/Baker PRIMNM and the Phoenix Islands Protected Area (PIPA). A scale bar indicates distances up to 200 nautical miles. A north arrow is located in the bottom right corner.
Site Name	Carondelet Reef
Expedition Coordinator(s)	Brian RC Kennedy, Nick Pawlenko
ROV Lead(s)	Karl McLetchie
Science Team Lead(s)	Amanda Demopoulos and Steven Auscavitch
General Area Descriptor	Phoenix Islands Protected Area
ROV Dive Name	
Cruise	EX-17-03

	Amanda Demopoulos	USGS	ademopoulos@usgs.gov	
	Amanda Netburn	NOAA OER	amanda.netburn@noaa.gov	
	Amy Baco-Taylor	Florida State University	abacotaylor@fsu.edu	
	Andrea Pierce	UnderWater World Guam	apierce@aquariumteam.com	
	Asako Matsumoto	Chiba Institute of Technology (Chitech),	amatsu@gorgonian.jp	
	Bruce Mundy	NOAA NMFS Pacific Islands Fisheries Science Center	bruce.mundy@noaa.gov	
	Christopher Kelley	University of Hawaii	ckelley@hawaii.edu	
	Erik Cordes	Temple University	ecordes@temple.edu	
	Jaymes Awbrey	University of Louisiana, Lafayette	jawbrey@louisiana.edu	
	Jill Bourque	US Geological Survey Wetland and Aquatic Research Center	jbourque@usgs.gov	
	Ken Sulak	USGS emeritus	ksulak@usgs.gov	
	Les Watling	University of Hawaii at Manoa	watling@hawaii.edu	
	Michael Parke	NOAA PIFSC	michael.parke@noaa.gov	
	Natalie Summers	University of Hawaii at Manoa	nsummers@hawaii.edu	
	Nicole Morgan	Florida State University	nmorgan@fsu.edu	
	Peter Auster	Mystic Aquarium & UConn	peter.auster@uconn.edu	
	Randi Rotjan	Boston University	rrotjan@bu.edu	

	Santiago Herrera	Lehigh University	sherrera@alum.mit.edu
	Scott France	University of Louisiana at Lafayette	france@louisiana.edu
	Steve Auscavitch	Temple University	steven.auscavitch@temple.edu
	Taylor Heyl	WHOI	theyl@whoi.edu
	Tina Molodtsova	P.P.Shirshov Institute of Oceanology RAS	tina.molodtsova@gmail.com
	Mike Ford	NOAA NMFS	michael.ford@noaa.gov
Purpose of the Dive	The goal of this dive is to acquire baseline information on deep sea habitats, seafloor geology, and biological communities in the Carondelet Reef area in the Phoenix Islands Protected Area (PIPA). This will be the first feature in the EX1703 expedition that will lie within the boundaries of PIPA. Deep-sea environments in PIPA are virtually unexplored leading to poor knowledge of biological resources protected by the MPA. This feature has a summit depth of <5m, capped by a submerged reef complex. The base lies at a depth of 5300m.		
Description of the Dive	The third dive of EX1703 was the most biologically diverse of all the dives thus far of the expedition. The dive track was on the southwest ridge of Carondelet Reef, the first dive for the cruise within the Phoenix Islands Protected Area (PIPA) and the first deep dive ever on Carondelet Reef. Our primary objective was to descend to 1840 m and traverse up a steep slope up to approximately 1630 m, then ascend along a more gradual ridge, up to 1587m. Throughout the dive, we planned to document corals, fishes, sponges and other invertebrates, while recording the dominant terrain type. We documented at least 32 species of corals, 21+ species of other invertebrates, including coral and sponge associates, and 5 species of fishes. The start of the dive, which traversed a vertical wall, was very speciose. Within the first 20 m of vertical water depth, we encountered at least 11 species of corals (<i>Paragorgia</i> sp. [collected specimen], 2 types of chrysogorgiids [some with eggs], 1 <i>Iridogorgia</i>, <i>Hemicorallium</i>?, 4 black corals [3 branched forms [<i>Trissopathes</i>?, <i>Stauropathes</i>?, <i>Bathypathes</i> sp.] and whips [<i>Stichopathes</i>?], 2 isidids), 2 sponges, and several associates (e.g., ophiuroids, crinoids,		

	amphipods, anemones, chirostylids, hydroids, barnacles). <i>Hemicorallium</i> is a known precious coral and the first one observed on the dives thus far. Several large bamboo colonies were observed attached to steep rock faces. Most of the fan corals were oriented perpendicular to the dominant current, which was flowing from NW to SE. As the dive continued upslope, we saw many of the same species of corals, plus a few new ones: <i>Bathypathes?</i> sp. (yellow), rock pens (<i>Anthoptilum</i>), <i>Umbellula</i>, <i>Pleurogorgia</i>, stoloniferans, <i>Iridogorgia magnispiralis</i>, <i>Anthomastus tahinodus</i>, acanthogorgiids, <i>Isidella</i> sp., and <i>Umbellapathes?</i>. Cup corals were the only scleractinians documented on the entire dive. Five species and 5 individuals of fish were observed throughout the dive: <i>Synaphobranchus</i> spp (2), Unknown Synaphobranchidae (1 sp.), batfish (Ogcocephalidae), and a bristlemouth (Gonostomatidae). The synaphobranchids were not observed on previous dives within American Samoa at similar depths. As we moved to gently sloped terrain (~1700 m water depth), there were more sedimented ledges and rock fissures filled with sediment. The substrate had several extremely long (>1 m) bamboo coral whips (Clade B1?). We also observed a large bamboo colony, possibly the largest ever observed, estimated to be 300 years old, with several tendrils that attached the skeleton to the substrate, securing its position on a steep rock ledge. We successfully sampled a similar isidid adjacent to the old one to confirm identification. Following this collection, we collected an unknown sponge from the family Euretidae with associates (crinoid and ophiuroids). Other sponges observed included hexactinellids: euplectellid, <i>Poliopogon</i> sp., <i>Amphidiscella</i> sp.?, and <i>Tretopleura</i> sp. The underside of rock ledges were often populated by small colonies of <i>Pleurogorgia</i>. Notable non-coral observations included predation by a seastar (<i>Circeaster pullus</i>) on a bamboo whip (Clade B1), homolid crab with an anemone clasped by its posterior limbs, a benthic siphonophore (Rhodaliidae?) that resembled a pipe organ, and an enormous (~17 cm) nemertean moving its head along the rocky seafloor. Our observations directly help the management of biological resources within PIPA, given nothing was previously known about the deep-sea fauna within the protected area. Future dives will enable us to clarify if the diversity and abundance of fauna present on Carondelet Reef is representative of the broader region. Images: big bamboo coral with tendrils, rock with lots of corals in view, seastar eating bamboo whip, batfish, and big nemertean.
--	--

Overall Map of the ROV Dive Area		Close-up Map of Main Dive Site	
			
Representative Photos of the Dive			
			
seastar (<i>Circeaster pullus</i>) feeding on a bamboo whip (Clade B1)		D2 looks at an unknown large big bamboo coral with tendrils	
Samples Collected			
Sample			
Sample ID	EX1703_20170310T201405_D2_DIVE03_SPEC01BIO		
Date (UTC)	20170310		
Time (UTC)	20:14:05		
Depth (m)	1841.11		
			

Temperature (°C)	2.37	
Field ID(s)	Paragorgia sp.	
Comments		
Sample		
Sample ID	EX1703_20170311T001717_D2_DIVE03_SPEC02BIO	
Date (UTC)	20170311	
Time (UTC)	00:17:17	
Depth (m)	1636.383	
Temperature (°C)	2.55	
Field ID(s)	Isididae	
Comments		
Sample		
Sample ID	EX1703_20170311T004803_D2_DIVE03_SPEC03BIO	
Date (UTC)	20170311	
Time (UTC)	00:48:03	
Depth (m)	1624.78	
Temperature (°C)	2.56	
Field ID(s)	Hexactinellida	
Comments		

Please direct inquiries to:

NOAA Office of Ocean Exploration & Research
1315 East-West Highway (SSMC3 10th Floor)
Silver Spring, MD 20910
(301) 734-1014